



CAUTION

**KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

ACTIVE CONSTITUENT:
750 g/kg CHLORSULFURON

CONTENTS:
20 g - 1 kg

GROUP	2	HERBICIDE
-------	----------	-----------

For the control of certain grass and broadleaved weeds in CleanCrop™ forage brassica crops as per the Directions for Use table.

SAFETY DIRECTIONS

May irritate the eyes. Avoid contact with eyes. When opening the container and preparing spray and using the prepared spray wear cotton overalls buttoned to the neck and wrist (or equivalent clothing). Wash hands after use.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

SAFETY DATA SHEET

For further information refer to the Safety Data Sheet available from www.fmccrop.com.au.

GENERAL INSTRUCTIONS

Telar™ is a selective systemic herbicide absorbed by the foliage and roots. It is rapidly translocated and acts by inhibiting acetolactate synthase (ALS), an enzyme necessary for root and shoot growth in sensitive plants.

Prior to using Telar™ careful consideration should be given to soil pH. As soil pH increases rate of breakdown decreases. Telar™ should not be used on soil pH 8.6 or higher as extended soil residual activity could adversely affect crop rotation options beyond normal intervals. Telar™ is intended for use on soils with a pH of 8.5 or lower.

Crops other than CleanCrop™ forage brassica varieties can be extremely sensitive to low concentrations of Telar™ in the soil. See Crop Rotation Recommendations, below.

Best weed control is obtained when rainfall or sprinkler irrigation wets the soil to a depth of 5 to 7.5 cm within 4 weeks of application.

RESISTANT WEEDS WARNING

GROUP	2	HERBICIDE
-------	----------	-----------

Telar™ is a member of the sulfonylurea group of herbicides. Telar™ has the inhibitor of the enzyme acetolactate synthase (ALS) mode of action. For weed resistance management Telar™ is a Group 2 herbicide.

Some naturally-occurring weed biotypes resistant to Telar™ and other Group 2 herbicides (Annual Ryegrass and some broadleaf weeds) are known to exist. These resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Telar™ or other Group 2 herbicides. Since the occurrence of resistant weeds is difficult to detect prior to use, FMC accepts no liability for any losses that may result from the failure of Telar™ to control resistant weeds.

Annual ryegrass biotypes resistant to Group 1 herbicides are often also resistant to Telar™. Before using Telar™ on a population resistant to Group 1 herbicides, have a resistance test conducted to ensure that it is still susceptible to Telar™.

To prevent, or at least minimise the risk of resistant weeds occurring, use Telar™ in tank mixes (if appropriate) and/or rotations with herbicides having different modes of action effective on the same weed species. Large numbers of healthy surviving weeds can be an indication that resistance is developing. Efforts should be taken to prevent seed set of these survivors.

DO NOT make more than two applications of a Group 2 herbicide to a crop, either pre-sowing incorporated by sowing (IBS)/post-sowing pre-emergence (PSPE) or post crop and weed emergence.

If the user suspects that a Group 2 herbicide resistant weed is present, Telar™ or other Group 2 herbicides should not be used. Strategies to minimise the risk of herbicide resistance are available. Consult your farm chemical supplier, consultant, or local FMC representative.

Telar™ Herbicide should be used as a part of an integrated weed management program which includes herbicides from other modes of action and non-chemical methods. Refer to the CropLife resistance management strategies, available from CropLife at www.croplife.org.au or contact your local FMC representative.

PRE-SOWING INCORPORATED BY SOWING

Avoid applying to dry sandy soils as rapid leaching may occur with early season rains (WA only).

When using Telar™ on light to medium soils pH 7 to 8.5, seek further advice from your FMC or DLF Seeds representative (SA only).

Conventionally Sown Crops – on soils of pH less than 7, apply a spray just before sowing or in conjunction with the sowing operation. On soils of pH of 7 or greater, it is not critical to time the spray just before sowing. Spray onto a non-ridged surface free of large clods. Use low profile 10 cm combine points for sowing. Sow at speeds of 10 kph or greater. Use light covering harrows at sowing. If applied to dry soil and sowing is to be delayed incorporate immediately after spraying to prevent loss by wind erosion.

Direct Drilled Crops – apply tank mixed with either Spray.Seed or glyphosate in accordance with manufacturer's label recommendations.

POST-SOWING OF THE CROP AND PRE-EMERGENCE OF THE WEEDS

Telar™ can be applied post sowing but pre-emergence of the CleanCrop™ crop. **DO NOT** apply Telar™ to emerging or the cotyledon stage of the CleanCrop™ forage brassica crop.

POST CROP AND WEED EMERGENCE

DO NOT apply Telar™ before the crop reaches the 4 leaves growth stage.

Where treatment is delayed or where weeds are not actively growing due to adverse conditions, results may be slow to appear and weeds may be only stunted or suppressed.

Apply after crop emergence at the 4-6 leaf stage of the Cleancrop™ forage brassica crop, when weeds are small and actively growing (Annual ryegrass no more than 3 leaves, refer to diagram in the Directions for Use table) Broadleaved weeds no more than 5 cm in height or diameter (for Black bindweed refer to specific recommendations) and before crop canopy closure.

GRAZING ADVICE

A 14-day withholding period is applicable for grazing Telar™ treated areas. Observing this withholding period after application will also optimise weed control.

CROP SAFETY

Application of Telar™ may result in transient yellowing or bleaching of Cleancrop™ forage brassica crops. Recovery of the crop from these transient symptoms will be dependent on the Cleancrop™ brassica variety and may take up to several weeks under cold climatic conditions.

DO NOT apply Telar™ to crops that are stressed by any cause (such as severe weather conditions, drought, waterlogging, frosts, excessive soil acidity or alkalinity, poor nutrient status, disease, nematode or insect damage) as enhanced and prolonged crop injury may result. When treatment is followed by severe stress (such as drought, prolonged cold, waterlogging or frosty conditions) crop yellowing and growth retardation may occur. Crops normally recover without loss in yield. Disease, nematode or insect damage following application may also result in crop injury.

The use of Telar™ on Cleancrop™ forage brassica varieties sown out of season may result in some yield suppression under stressed conditions (e.g. Aluminium/Manganese toxicity, waterlogging, frosts, etc.). Contact your local FMC or DLF Seeds representative on the seasonal suitability of the Cleancrop™ forage brassica of choice for your district.

Use of Telar™ in furrow or flood-irrigated areas

Pre-sowing irrigation should be no closer than 10 days before application of Telar™.

DO NOT furrow or flood-irrigate areas treated with Telar™ if water logging or other stress factors will be induced.

DO NOT apply Telar™ to light, sandy ridges low in organic matter, as crop injury may occur.

Consult your local FMC or DLF Seeds Representative for latest information relating to variety tolerance to Telar™.

CROP ROTATION RECOMMENDATIONS

Land previously treated with Telar™ should not be rotated to other crops other than those listed in the following tables.

Tolerance of other crops (grown through to maturity) should be determined on a small scale before sowing into larger areas. DO NOT replant any other crop, other than those listed in the Tables below, without first consulting your local FMC or DLF Seeds Representative.

The Telar™ treated area may be re-planted to any of the specified crops after the interval indicated in the following tables:

NB - THE TABLE BELOW APPLIES TO ALL STATES

MINIMUM RECROPPING INTERVAL (MONTHS AFTER APPLICATION)						
Soil pH*	0	3	6	9	12	18
6.5 or less	Triticale Wheat Cleancrop™ forage brassica	Cereal rye Canola <i>(Designated Imidazolinone herbicide tolerant Canola varieties only)</i>	Oats	Barley	Canola/Rapeseed Subterranean clover** Faba beans Field pea Linseed Lucerne Lupins Medics** Safflower	Maize Sorghum Soybeans Sunflower

NB - THE TABLES BELOW APPLY TO Qld, SA, WA & Tas ONLY

MINIMUM RECROPPING INTERVAL (MONTHS AFTER APPLICATION)						
	0	3	9	15	18	22
Rainfall requirement				Minimum 700 mm		
Soil pH* 6.6 - 7.5	Triticale Wheat CleanCrop™ forage brassica	Cereal Rye Canola <i>(Designated Imidazolinone herbicide tolerant Canola varieties only)</i>	Barley Oats	Japanese millet Maize Panicum millet Sorghum Sunflower White French millet	Cotton Soybeans	Canola/Rapeseed Faba Beans Field Pea Linseed Medics** Rapeseed Safflower Subterranean clover**

MINIMUM RECROPPING INTERVAL (MONTHS AFTER APPLICATION)					
	0	3	15	18	24 months or longer
Rainfall Requirement	Minimum 700 mm				
Soil pH* 7.6 - 8.5	Triticale Wheat CleanCrop™ forage brassica	Canola <i>(Designated Imidazolinone herbicide tolerant Canola varieties only)</i>	Japanese millet Maize Panicum millet Sorghum Sunflowers White French millet	Barley Oats Cereal rye	Rotate to crops other than Cereals and Cleancrop™ forage brassica (such as listed above) only if field test strip of the planned rotational crop has been successfully grown through to maturity in the previous season
8.6 and above	Telar™ is not recommended for use on soils of pH 8.6 & above.				

NB - THE TABLES BELOW APPLY TO NSW & Vic ONLY

MINIMUM RECROPPING INTERVAL (MONTHS AFTER APPLICATION)					
Soil pH*	0	3	9	22	26
6.6 - 7.5	Triticale Wheat CleanCrop™ forage brassica	Cereal rye Canola <i>(Designated Imidazolinone herbicide tolerant Canola varieties only)</i>	Barley Oats	Canola/Rapeseed Subterranean Clover** Faba beans Field peas Linseed Lucerne Lupins Medics** Safflower	Maize Sorghum Soybeans Sunflower

MINIMUM RECROPPING INTERVAL (MONTHS AFTER APPLICATION)				
Soil pH*	0	3	18	
7.6 - 8.5	Triticale Wheat CleanCrop™ forage brassica	Canola <i>(Designated Imidazolinone herbicide tolerant Canola varieties only)</i>	Barley, Cereal rye, Oats	Rotate to crops other than Cereals and CleanCrop™ forage brassica (such as listed above) only if field test strip of the planned rotational crop has been successfully grown through to maturity in the previous season
8.6 and above	Telar™ is not recommended for use on soils of pH 8.6 & above.			

* Soil pH is to be determined by Laboratory analysis using the 1:5 soil: water suspension method.

** Includes natural regeneration of subterranean clover and medics.

SPRAY PREPARATION

Telar™ is a water dispersible granule formulation to be mixed with water and be applied as a spray. Partially fill the spray tank with water. Using chemical resistant gloves, measure the amount of Telar™ required for the area to be sprayed using the graduated measuring cylinder provided. Before spraying calibrate equipment to determine the quantity of water necessary to uniformly cover the measured area to be treated. Add the correct amount of Telar™ to the spray tank with the agitation system engaged. Top up to the correct volume with water. **THE PRODUCT MUST BE KEPT IN SUSPENSION AT ALL TIMES BY CONTINUOUS AGITATION.**

In tank mixes Telar™ must be in suspension before adding the companion herbicide or surfactant.

For application of Telar™ in liquid fertiliser, slurry the Telar™ in water, then thoroughly mix the slurry into the liquid fertiliser. Check compatibility before mixing Telar™ in fertiliser solutions.

When mixing Telar™ in a tank with other products, the following mixing sequence should be followed:

1. Water conditioning agents;
2. Water dispersible granules (WG) (e.g. Telar™ Herbicide);
3. Wettable powders (WP);
4. Flowables or suspension concentrates (SC);
5. Water-soluble concentrates (e.g. glyphosate);
6. Emulsifiable concentrates (EC) – see Note below;
7. Surfactant and oils (e.g. BS1000#) – note for use with post-emergence applications of Telar™ **ONLY**.

INCOMPATIBILITY

Note: Telar™ when applied post-emergence with EC formulations will cause severe crop damage. **DO NOT** tank mix Telar™ with EC formulations for post-emergence applications.

Where prepared spray solutions have been allowed to stand, thoroughly re-agitate before using.

COMPATIBILITY

The use of spraying oils required with some insecticides may cause increased phytotoxicity to the crop.

For information on compatibility with specific product formulations refer to the comprehensive compatibility guide for Telar™ either online (www.fmccrop.com.au) or seek advice from your local FMC or DLF Seeds representative.

USE OF SURFACTANT/WETTING AGENT

For post-emergence application, always add a non-ionic surfactant (1000 g ai/L), e.g. BS1000# at 100 mL/100 L of final spray volume (0.1 % v/v).

The use of spraying oils may cause increased phytotoxicity to the crop.

NOTE: DO NOT add a surfactant/wetting agent when Telar™ is tank mixed with another product which already has a surfactant/wetting agent in the formulation.

GROUND SPRAYING EQUIPMENT

Use a boom spray properly calibrated to a constant speed and rate of delivery to ensure thorough coverage and a uniform spray pattern.

Avoid overlapping and shut off spray booms while starting, turning slowing or stopping as injury to the crop may result.

A minimum spray volume of 60 L water/hectare must be applied. Spray volumes of 60 - 200 L water/ha may be used.

AERIAL APPLICATION

Apply a minimum of 30 L/ha water (fixed wing aircraft) and 50 L/ha (helicopter). Avoid spraying in still conditions and in winds likely to cause drift onto adjacent sensitive crops. Avoid spraying where drift can go onto areas likely to be sown to sensitive crops - see Crop Rotation Recommendations. Turn off spray boom whilst passing over creeks and dams.

SPRAYER CLEANUP

To avoid subsequent injury to crops other than wheat, triticale, barley, oats, cereal rye or Cleancrop™ forage brassica, immediately after spraying thoroughly remove all traces of Telar™ from mixing and spray equipment as described below. Ensure that the following operation is carried out **in an area** that is clear of waterways, desirable vegetation and tree roots, and preferably in an area where drainings can be contained.

1. Drain tank, then flush tank, boom and hoses with clean water for a minimum of 10 minutes.
2. Fill the tank with clean water then add 300 mL household chlorine bleach (containing 4 % chlorine) per 100 L of water. Flush through boom and hoses then allow to sit for 15 minutes with agitation engaged, then drain.
3. Repeat Step 2.
4. Nozzles and screens should be removed and cleaned separately. To remove traces of chlorine bleach, rinse the tank thoroughly with clean water and flush through hoses and boom.

CAUTION:

DO NOT use chlorine bleach with ammonia. All traces of liquid fertiliser containing ammonia, ammonium nitrate or ammonium sulphate must be rinsed with water from the mixing and application equipment before adding chlorine bleach solution. Failure to do so will release a gas with a musty chlorine odour that can cause eye, nose, throat and lung irritation.

DO NOT clean equipment in an enclosed area.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

Injury to or loss of desirable trees or vegetation may result from failure to observe the following:

DO NOT apply or drain or flush equipment on or near desirable trees or other plants or on areas where their roots may extend or in locations where the chemical may be washed or moved into contact with their roots.

DO NOT allow drainage water from irrigated areas treated with Telar™ to enter regional drains, local waterways or river systems.

DO NOT apply drainage water from irrigated areas treated with Telar™ onto sensitive crops.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Very toxic to aquatic life. **DO NOT** contaminate wetlands or watercourses with this product or used containers.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area.

DO NOT store for prolonged periods in direct sunlight. Store in a locked room or place away from children, animals, food and feedstuffs, seed in open packaging and fertilisers.

DO NOT re-use container. Triple-rinse containers before disposal. Add rinsings to spray tank. **DO NOT** dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging to an approved waste management facility.

If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with local, state or territory government regulations. **DO NOT** burn empty containers or product.

NOTICE TO BUYER

To the extent permitted by the Competition and Consumer Act (2010) or any relevant legislation of any State or Territory (the "Legislation") all conditions and warranties and statutory or other rights of action, whether arising in contract or tort or whether due to the negligence of FMC or Seller, which buyer or any other user may have against FMC or Seller are hereby excluded provided however that any rights of the buyer pursuant to non-excludable conditions or warranties of the Legislation are expressly preserved. FMC hereby gives notice to buyer and other users that to the extent permitted by the Legislation it will not accept responsibility for any indirect or consequential loss of whatsoever nature arising from the storage, handling or use of this Product. Where permitted by the Legislation FMC's liability shall in all circumstances be limited to the replacement of the product, or a refund of the purchase price paid therefor.

The Product must be used and applied strictly in accordance with the label instructions and other directions for use. It is impossible to eliminate all risks associated with the use of this product. Such risks may arise from factors such as weather conditions, soil factors, off target movement, unconventional technique, presence of other materials, the manner of use or application, or other unknown factors, all of which are beyond the control of FMC or the Seller. Buyer accepts these risks.

APVMA Approval Number: 90141/135834

FMC, the FMC logo and Telar are trademarks of FMC Corporation and/or an affiliate.

©2023 FMC Corporation. All rights reserved.

CleanCrop™ is a trademark of DLF Seeds.

Non FMC Trademark

FMC Australasia Pty Ltd.
A.B.N. 45 095 326 891
Level 2, Building B
12 Julius Ave
North Ryde NSW 2113
Phone: 1800 066 355
www.fmccrop.com.au

SPECIALIST ADVICE IN EMERGENCY ONLY
1800 033 111 ALL HOURS – AUSTRALIA WIDE

Additional statements required by Globally Harmonised Systems for classification and labelling of chemicals (GHS) and Safe Work Australia: very toxic to aquatic life with long lasting effects.

Collect spillage. Not combustible. Use extinguishing media suited to burning materials. Suitable extinguishing media: carbon dioxide, dry chemical, water spray, foam. Do not use high volume water jet.

STORAGE: Protect from sunlight. Store in a dry place. Store in a closed container. Store in a well-ventilated place. Keep cool. Dispose of contents and containers as specified on the registered label.

RESTRAINTS:

DO NOT apply if heavy rains or storms are forecast within 3 days.

DO NOT irrigate to the point of field runoff for at least 3 days after application.

DO NOT allow drainage water from irrigated areas treated with Telar™ to enter regional drains, local waterways or river systems.

DO NOT store a suspension of Telar™ for more than 2 days otherwise significant breakdown will occur.

DO NOT store tank mixes of Telar™ with other materials for more than 24 hours as product breakdown may occur.

DO NOT spray for post emergent weed control if rainfall is expected within two (2) hours.

DO NOT apply to emerging or cotyledon stage or before the four (4) leaf stage of the CleanCrop™ forage brassica crops.

DO NOT apply Telar™ to CleanCrop™ forage brassica crops which are undersown with other species including other brassicas, clover, lucerne and grass species.

SPRAY DRIFT RESTRAINTS

Specific definitions for terms used in this section of the label can be found at apvma.gov.au/spraydrift.

DO NOT allow bystanders to come into contact with the spray cloud.

DO NOT apply in a manner that may cause an unacceptable impact to native vegetation, agricultural crops, landscaped gardens and aquaculture production, or cause contamination of plant or livestock commodities, outside the application site from spray drift. Wherever possible, correctly use application equipment designed to reduce spray drift and apply when the wind direction is away from these sensitive areas.

DO NOT apply unless the wind speed is between 3 and 20 kilometres per hour at the application site during the time of application.

DO NOT apply if there are hazardous surface temperature inversion conditions present at the application site during the time of application. Surface temperature inversion conditions exist most evenings one to two hours before sunset and persist until one to two hours after sunrise.

DO NOT apply by a boom sprayer unless the following requirements are met:

- Spray droplets are not smaller than a **COARSE** spray droplet size category.

DO NOT apply by aircraft unless the following requirements are met:

- Spray droplets not smaller than a **COARSE** spray droplet size category.
- Maximum release height above the target canopy of 3 metres or 25 per cent of wingspan or 25 per cent of rotor diameter, whichever is the greatest.

DIRECTIONS FOR USE

USE METHOD – PRE-SOWING INCORPORATED BY SOWING or POST-SOWING PRE-EMERGENCE

Grass weeds – All States

CROP	TARGET SPECIES	RATE g/ha	CRITICAL COMMENTS
CleanCrop™ varieties of forage brassica	Annual (Wimmera) ryegrass (<i>Lolium rigidum</i>)	20	NOTE:- Refer to General instructions on page 1 for optimum application timing and conditions.
	Brome grass (<i>Bromus</i> spp) (Suppression only)		Gives suppression only if populations are 20 plants/m ² or less.
	Paradoxa grass (<i>Phalaris paradoxa</i>)		Use only on soils pH >7.5. Apply to dry soil before the sowing rain. Mechanical incorporation before the sowing rains is not necessary.

USE METHOD – PRE-SOWING INCORPORATED BY SOWING or POST-SOWING PRE-EMERGENCE

Broadleaf weeds – All States

CROP	TARGET SPECIES	RATE g/ha	CRITICAL COMMENTS
CleanCrop™ varieties of forage brassica	African Turnip weed (<i>Sisymbrium thellungii</i>)	20	Apply to dry soil before the sowing rain. Mechanical incorporation before the sowing rains is not necessary.
	Amsinckia/Yellow Burrweed (<i>Amsinckia</i> spp.)		
	Ball Mustard (<i>Neslia puniculata</i>)		
	Black Bindweed/Climbing Buckwheat (<i>Fallopia convolvulus</i>)		
	Cape Tulip (<i>Homeria</i> spp.)		


CROP	TARGET SPECIES	RATE g/ha	CRITICAL COMMENTS
CleanCrop™ varieties of forage brassica <i>continued</i>	Capeweed (<i>Arctotheca calendula</i>)	20	On acid soils pH 5.5 or less this product will give a shorter period of control in wet years.
	Charlock (<i>Sinapis arvensis</i>)		
	Common Iceplant (<i>Mesembryanthemum crystallinum</i>)		
	Corn Gromwell/Sheepweed/White Ironweed (<i>Buglossoides arvensis</i>)		
	Deadnettle (<i>Lamium amplexicaule</i>)		
	Docks (<i>Rumex</i> spp.)		
	Fat-hen (<i>Chenopodium album</i>) (Suppression only)		
	Fumitory (<i>Fumaria</i> spp.) (Suppression only)		
	Guildford grass/Onion grass (<i>Romulea rosea</i>)		
	Indian Hedge Mustard (<i>Sisymbrium orientale</i>)		
	King Island Melilot (<i>Melilotus indicus</i>)		
	Lincoln weed (<i>Diplotaxis tenuifolia</i>)		
	Loosestrife (<i>Lythrum</i> spp.)		
	Mintweed (<i>Salvia reflexa</i>)		
	Mouse-ear Chickweed (<i>Cerastium</i> spp.)		
	New Zealand Spinach (<i>Tetragonia tetragonoides</i>)		
	Paterson's Curse/Salvation Jane (<i>Echium plantagineum</i>)		
	Pimpernels (<i>Anagallis arvensis</i>)		
	Prickly Lettuce/Whip Thistle (<i>Lactuca serriola</i>)		
	Rough Poppy (<i>Papaver hybridum</i>)		
	Saffron Thistle (<i>Carthamus lanatus</i>) (Suppression only)		
	Saltbush (<i>Atriplex muelleri</i>)		
	Shepherd's Purse (<i>Capsella bursa-pastoris</i>)		
	Slender Celery (<i>Apium leptophyllum</i>)		
	Slender Thistle (<i>Carduus tenuiflorus</i>)		
	Soursob (<i>Oxalis pes-caprae</i>)		
Spear Thistle (<i>Cirsium vulgare</i>)			

CROP	TARGET SPECIES	RATE g/ha	CRITICAL COMMENTS
CleanCrop™ varieties of forage brassica <i>continued</i>	Stemless Thistle (<i>Onopordum acaulon</i>)	20	
	Storksbill/Wild Geranium (<i>Erodium</i> spp.)		
	Three Cornered Jack(s) / Doublegee/ Spiny Emex (<i>Emex australis</i>)		
	Tree Hogweed (<i>Polygonum patulum</i>)		
	Turnip weed (<i>Rapistrum rugosum</i>)		
	Wireweed/Hogweed (<i>Polygonum aviculare</i>)		
	Wild Turnip (<i>Brassica tournefortii</i>)		

USE METHOD – POST CROP (4 true leaf stage onwards) AND WEED EMERGENCE

Pre-emergence application (IBS or PSPE) of Telar™ Herbicide may be followed by a second application of Telar® Herbicide or a suitable post-emergence herbicide to control any newly emerged weeds.

Grass weeds – All States

CROP	TARGET SPECIES	RATE g/ha	CRITICAL COMMENTS
CleanCrop™ varieties of forage brassica	Annual (Wimmera) ryegrass (<i>Lolium rigidum</i>) (Control of Annual (Wimmera) ryegrass under low to moderate weed pressure. Suppression only under high weed pressure.)	20	 <p>1 leaf 2 leaf 3 leaf 4 leaf Too late to spray</p> <p>Apply no later than the 3-leaf stage of Annual ryegrass.</p> <p>Application of Telar™ to Annual ryegrass with water volumes less than 60 L/ha may result in reduced efficacy.</p>

USE METHOD - POST CROP (4 true leaf stage onwards) AND WEED EMERGENCE

Pre-emergence application (IBS or PSPE) of Telar™ Herbicide may be followed by a second application of Telar™ Herbicide or a suitable post-emergence herbicide to control any newly emerged weeds.

Broadleaf weeds – All States

CROP	TARGET SPECIES	RATE g/ha	CRITICAL COMMENTS
CleanCrop™ varieties of forage brassica	African Turnip weed (<i>Sisymbrium thellungii</i>)	20	Apply at cotyledon to 4 leaf stage.
	Amsinckia/ Yellow Burrweed (<i>Amsinckia</i> spp.)		
	Ball Mustard (<i>Neslia puniculata</i>)		
	Bifora/ Carrot weed (<i>Cotula australis</i>) (Suppression only)		
	Black Bindweed/Climbing Buckwheat (<i>Fallopia convolvulus</i>)		Apply at cotyledon to 2 leaf stage of weed.
	Cape Tulip (<i>Homeria</i> spp.)		
	Charlock (<i>Sinapis arvensis</i>)		
	Corn Gromwell/ Sheepweed/White Ironweed (<i>Buglossoides arvensis</i>)		Apply at cotyledon to 2 leaf stage. If applied at a later stage only suppression will occur.
	Deadnettle (<i>Lamium amplexicaule</i>)		
	Docks (<i>Rumex</i> spp.)		

CROP	TARGET SPECIES	RATE g/ha	CRITICAL COMMENTS
	Fat-hen (<i>Chenopodium album</i>)		
CleanCrop™ varieties of forage brassica <i>continued</i>	Fumitory, Denseflower (<i>Fumaria densiflora</i>)	20	Apply at cotyledon to 2 leaf stage.
	Guildford grass/Onion grass (<i>Romulea rosea</i>)		
	Hoary Cress (<i>Cardaria draba</i>)		Apply when plants are fully emerged.
	Lincoln weed (<i>Diplotaxis tenuifolia</i>)		
	Matricaria (<i>Matricaria matricarioides</i>)		
	Mintweed (<i>Salvia reflexa</i>)		Apply at cotyledon to 4 leaf stage.
	Mouse-ear Chickweed (<i>Cerastium</i> spp.)		
	Mustards (<i>Sisymbrium</i> spp.)		
	New Zealand Spinach (<i>Tetragonia tetragonoides</i>)		
	Paterson's Curse/Salvation Jane (<i>Echium plantagineum</i>)		
	Pimpernels (<i>Anagallis arvensis</i>)		
	Prickly Lettuce/Whip Thistle (<i>Lactuca scariola</i>)		
	Rough Poppy (<i>Papaver hybridum</i>)		
	Saltbush (<i>Atriplex muelleri</i>)		Apply at cotyledon to 4 leaf stage.
	Shepherd's Purse (<i>Capsella bursa-pastoris</i>)		
	Slender Celery (<i>Apium leptophyllum</i>)		Apply at cotyledon to 4 leaf stage.
	Soursob (<i>Oxalis pes-caprae</i>)		Apply when the majority of Soursobs have emerged.
	Spear Thistle (<i>Cirsium vulgare</i>)		A follow-up spray with a suitable herbicide may be necessary to control subsequent germinations.
	Stagger weed (<i>Stachys arvensis</i>)		
	Stemless Thistle (<i>Onopordum acaulon</i>) (Suppression only)		
	Storksbill/Wild Geranium (<i>Erodium</i> spp.)		
	Tree Hogweed (<i>Polygonum patulum</i>)		
	Turnip weed (<i>Rapistrum rugosum</i>)		
Wild Radish (<i>Raphanus raphanistrum</i>)	A follow-up spray with a suitable herbicide may be necessary to control subsequent germinations.		
Wild Turnip (<i>Brassica tournefortii</i>)			
Wireweed/Hogweed (<i>Polygonum aviculare</i>)			

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

GRAZING WITHHOLDING PERIOD

CLEANCROP™ FORAGE BRASSICAS: DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 14 DAYS AFTER APPLICATION